

Seasonal and Spatial Variation of Epilithic Algal Community in Batlama Stream (Giresun, Turkey)

Faruk Maraşlıoğlu¹, Elif Neyran Soylu², Sibel Altürk Karaca²

¹ Hitit University, Department of Environmental Protection Technologies, Çorum, TURKEY

² Giresun University, Department of Biology, Giresun, TURKEY

ABSTRACT

Spatio-temporal changes in taxonomic composition and structure of epilithic algal community with some physicochemical features of the Batlama stream were assessed between June 2013 and May 2014 at the intertidal zone of the Batlama Stream, Giresun, Turkey. A total of 90 taxa were identified belonging to division of Bacillariophyta (80 taxa), Euglenophyta (3 taxa), Cyanobacteria (3 taxa), Charophyta (2 taxa) and Chlorophyta (2 taxa) on epilithic algae of Batlama Stream. *Encyonema minutum*, *Ulnaria ulna*, *Cocconeis placentula*, and *Navicula cryptocephala* species were the most abundant taxa among all samples of the four stations throughout the study.

The Shannon diversity index (H') values varied in the range from 0.5 to 1.2 and correlated with both species richness and relative species abundance (evenness). The results of the diversity analysis and the counting did not exactly match up with each other. Chlorophyll-a values of the stream and trophic classification (TDI, BDI) based on epilithic diatoms ascribed Batlama Stream to the mezotrophic range. Also, according to the calculated pollution index (PTI, S) values, all stations of the Batlama Stream were moderately polluted (β -mesosaprobity).

Keywords:

Stream; Epilithic algae; Seasonal variation; Diversity; Diatom index.

INTRODUCTION

Diatoms, besides the role of environmental condition indicator, also determine the water quality and are as interesting and intriguing phytoplanktons, as snowflakes. They took a primary role in various studies, in terms of oxygen, nutritional value and pollution indicator. The main primary producers in undisturbed and shaded zones of the streams are epilithic algae and play major roles in controlling energy flow of food webs in stream ecosystems [1, 2]. However, the overgrowth of epilithic algae may deteriorate water quality [3]. The production and dynamics of an epilithic algal composition in stream ecosystems are also largely influenced by physical variables such as flow rate, light and water temperature [4, 5]. The variation of epilithic algal biomass may indirectly affect the distribution of the Batlama stream. Water quality variables (electrical conductivity, pH, total dissolved solids, chemical oxygen demand, biochemical oxygen demand, and nutrients such as phosphorus and nitrogen from the surrounding lands) also play major roles in regulating the production rate and species com-

position of epilithic algae in streams [6]. However, in complex stream ecosystems, the dynamics of epilithic algal biomass may nonlinearly interact with the combination of abiotic and biotic factors. Therefore, some index varieties may help to describe the nonlinear relationships between epilithic algal biomass and aforementioned factors.

In Turkey, studies of diatom were began by Yıldız in Meram stream [7] and followed by other researchers which mostly concentrated on various physicochemical factors affecting the development of diatoms in addition to their flora in central anatolian [9, 10, 11, 12, 13, 14, 15, 16]. In recent years, some topics that determine trophic state and water quality level of a stream monitoring by means of diatom indices based on the variety and density of the diatoms in water or on the substrat have become popular [17, 18, 19].

The TDI was designed with the practical needs of busy water industry biologists in mind and, as a result,

Article History:

Received: 2017/01/06

Accepted: 2017/02/18

Online: 2017/06/30

Correspondence to: Faruk Maraşlıoğlu,
Hitit University, Department of
Environmental Protection Technologies,
Çorum, TURKEY

Tel: +90 (364) 223-8000-3376

Fax: +90 (364) 223-8004

E-Mail: farukmaraslioglu@hitit.edu.tr

was designed to be robust and easily learnt. It uses the weighted average equation of Zelinka and Marvan [20] to interpret benthic diatom community structure in terms of the nutrient concentrations in the river.

Epilithic algal samples obtained in the present study comprised of mixed forms growing on silicate-rich stones and phytoplankton forms commonly living in littoral and pelagic zones of freshwaters, however, in this context this all community was termed as epilithon. The aims of the present study were to describe the species composition of diatom assemblages, to determine which environmental factors explain the composition of epilithic diatoms and to assess the response of common diatom taxa to these environmental variables..

MATERIALS AND METHODS

Samples analysed in this work were collected monthly between June 2013 and May 2014 from four stations (Fig. 1). Stones collected from the stations in the stream were brushed clean of algae and the samples fixed the formaldehyde solution (4%). The epilithic diatoms were then placed on permanent slides which had been prepared according to Round [21]. At least 300 diatom valves per slide were counted at 400x magnification on Olympus BX51 microscope. In the evaluations, the average of three countings from the station was used [22]. Species identifications were based primarily on the John *et al.* [23], Krammer and Lange-Bertalot [24, 25, 26, 27]. In addition, current systematic status of algae taxa is checked by way of AlgaeBase databases [28] and the author names are given in abbreviated form according to Brummit and Powel [29].

The Shannon diversity index, the evenness and species

richness were applied on epilithic flora at different stations. The software used was PRIMER version 5.0 from Plymouth Marine Laboratory for Shannon-Wiener index [30]. Trophic diatom index (TDI) and Biological Diatom Index (BDI) were calculated by using the formula [31] and "IBD Calculate With Excel" programme [32], respectively. Also, the saprobic index (S) was evaluated based on the formula proposed by Pantle and Buck [33] and the diatom pollution tolerance index (PTI) was calculated based on the formula [34]. Water temperature (°C), pH, and dissolved oxygen concentration (mg/l) were measured in situ with WTW multi 340i/SET model pH meter. After filtering through GF/C and extracting in cold 90% acetone, chlorophyll-a (chl-a) was determined according to the equations of Strickland and Parsons [35].

RESULTS AND DISCUSSION

A total of 90 taxa belonging to division of Bacillariophyta (80), Euglenophyta (3), Cyanobacteria (3), Charophyta (2) and Chlorophyta (2) were identified in this study (Table 1). Bacillariophyta members were rich in species diversity and intensity in the algal flora of Batlama Stream (Fig. 2).

Encyonema minutum, *Ulnaria ulna*, *Cocconeis placentula*, and *Navicula cryptocephala* were predominantly found species in all sampling stations in the study area (Fig. 3). Sienkiewicz [36] stated that *U. ulna* have wide trophic tolerance and can live in oligo-eutrophic water. So, to be dominant of *U. ulna* throughout the study period point to a mesotrophic environment. *Navicula cryptocephala* and *Ulnaria ulna* are known that broad to distribution in Turkey [37, 38].

Total organisms decreased their numbers during the months of high rainfall. Sivacı and Dere [39] reached also a



Figure 1. Locations of the sampling stations in Batlama Stream

Table 1. The list of common taxa found in all four stations of the Batlama Stream

| Division: Bacillariophyta | |
|--|--|
| <i>Amphora ovalis</i> (Kütz.) Kütz. | <i>Melosira varians</i> C. Agardh |
| <i>Closterium navicula</i> (Bréb.) Lütkem. | <i>Navicula cryptocephala</i> Kütz. |
| <i>Cocconeis placentula</i> Ehrenb. | <i>Navicula lanceolata</i> (C. Agardh) Kütz. |
| <i>Cymatopleura solea</i> (Bréb.) W. Sm. | <i>Navicula menisculus</i> Schum. |
| <i>Cymbella affinis</i> Kütz. | <i>Navicula rhynchocephala</i> Kütz. |
| <i>Cymbella helvetica</i> Kütz. | <i>Navicula tripunctata</i> (O.F.Müller) Bory |
| <i>Cymbella tumida</i> (Bréb.) Van Heurck | <i>Navicula viridula</i> var. <i>linearis</i> Hustedt |
| <i>Diatoma vulgare</i> Bory | <i>Nitzschia acicularis</i> (Kütz.) W. Sm. |
| <i>Didymosphenia geminata</i> (Lyngb.) M.Schmidt | <i>Nitzschia palea</i> Kütz. |
| <i>Encyonema caespitosum</i> Kütz. | <i>Rhoicosphenia abbreviata</i> (C.Agardh) Lange-Bert. |
| <i>Encyonema minutum</i> (Hilse) D.G.Mann | <i>Surirella minuta</i> Bréb. |
| <i>Encyonema silesiacum</i> (Bleisch) D.G.Mann | <i>Ulnaria ulna</i> (Nitzsch) Compère |
| <i>Eunotia minor</i> (Kütz.) Grunov | Division: Chlorophyta |
| <i>Fragilaria capucina</i> Desmaz. | <i>Botryococcus braunii</i> Kütz. |
| <i>Gomphonema augur</i> Ehrenb. | Division: Cyanobacteria |
| <i>Gomphonema minutum</i> (C. Agardh) C. Agardh | <i>Oscillatoria tenuis</i> C. Agardh |
| <i>Gomphonema olivaceum</i> (Hornem.) Bréb. | Division: Euglenophyta |
| <i>Gyrosigma scalproides</i> (Rabenh.) Cleve | <i>Euglena gracilis</i> G.A.Klebs |
| <i>Hannaea arcus</i> (Ehrenb.) R.M.Patrick | <i>Euglena viridis</i> (O.F. Müll.) Ehrenb. |




Figure 2. % Distribution of Epilithic Algal Composition in Batlama Stream




Figure 3. Seasonal variations of predominant species in the epilithon of the Batlama stream

very similar conclusion during their study. Species diversity increased in the months of which the water temperature rise. Yavuz and Çetin [40] had recorded that relationship between seasonal variation of algae and water temperature.

The all parameters are high during the study period except winter. Water temperature ranged between 5.2 and

24.3°C in the stream. The development of epilithic algae in the stream affected by water temperature. The epilithic algal density was lower in some months except February and November than the other months when the temperature was low. In the stream throughout the research period, whereas the maximum oxygen level was 14 mg/l, the minimum level was 2 mg/l. The solubility of oxygen in the water decreases with the increase in temperature [41]. The density of oxygen in a natural oligotrophic freshwater having a temperature around 20°C is about 10 mg/l [42]. The average pH value of 7.8 indicates that the stream is slightly alkaline. The abundance of Encyonema minutum, Ulnaria ulna and Navicula cryptocephala in present study supports the Round's idea [43] that these species are widespread in alkaline waters. When total cell numbers were found to be high in autumn and winter seasons the pH values also reached similarly high values in these months in the study area. This result did not support the Vinebrooke's view [44] that algal growth decreases with pH increase.

The amount of chlorophyll-a were calculated monthly and chlorophyll-a showed usually similar values at the sampling stations. Chl-a ranged from 0.2 to 15.2 mg/m³ in Batlama stream. According to Moses [45] has stated that running water with chlorophyll-a values between 10 and 30 mg/m³ was mesotrophic. In term of the values, Batlama stream has a mesotrophic character.

The seasonal variation of Shannon diversity and even-




Figure 4. Shannon diversity, evenness, and species richness of the Batlama stream

ness did not vary greatly during the study period. While the highest values of the H' during the year was usually recorded in October, the lowest value was recorded in April. Recording the low value of Shannon diversity in April may be due to heavy rain mostly seen in Giresun in this month of the spring. In our study it was found that species diversity decreased in the spring and confirmed this with the Shannon diversity index. Species evenness index varied from 0.7–0.8. The lower species evenness values were often associated with the rainy season. Approaching the zero value of relative species abundance (evenness) indicates dominance of organism linked to single-species that *Encyonema minutum* and *Coconeis placentula* comprised of more than 80% of the all organisms in April 2014. Recording a decrease of H' and J' values with being the result of the excessive increase of these species indicates that Batlama stream is under pressure in these months of the year. The seasonal variations of species richness presented a decrease especially in cold period, while the highest values of species richness were recorded in autumn period (Fig. 4).

BDI and TDI indices provide information about trophic states of the lotic ecosystems. In the study, taxa with an mean relative abundance (%) less than 5 throughout the year were not included in the index calculation. Accordingly, 22 diatom taxa were used in St.1 and 19 diatom taxa in the other three stations to calculate the diatom indice scores of the four stations to determine the trophic status and water

quality of the Batlama Stream. While the lowest value of the TDI was recorded in station 3 as 49.8, the station has the lowest BDI value was station 4. Both values of the indices indicate oligo-mesotrophic features. TDI and BDI values of the four stations ranged between 50–55 and 14.7–15.4 respectively (Table 2). However, mean values of TDI (53.5) and BDI (15.1) indicated that the trophic structure of Batlama stream is mesotrophic. Similarly, according to PTI and S values, the studied site was moderately polluted (β -mesosaprobity). The present results coincide with the conclusions of Köster and Hübener [46], who suggested that the best method for monitoring freshwater by diatoms is the combined application of saprobity and trophic indices.

CONCLUSION

Agriculture, livestock activities and domestic waste water into the Batlama stream, are factors that increase of water pollution. The factors are decreased of density of epilithic algae. The present study results demonstrated that epilithic algal composition is distributed gradients of environmental factors. Temperature and rain are more important factor in explaining the seasonal variation of epilithic algae of Batlama stream. More research is needed to analyse the causes of the observed variation in the nutrient status and biomass of benthic algae in Batlama stream.

According to the diversity values, one can conclude that Batlama stream is on eutrophic level, however, the species

Table 2. Diatom indice values of the four stations in Batlama Stream derived from TDI, BDI, PTI and S

| | kind of index | Station 1 | Station 2 | Station 3 | Station 4 | Average |
|-----------------|---------------|-----------|-----------|-----------|-----------|---------|
| Trophic index | TDI | 53.8 | 55.3 | 49.8 | 55.3 | 53.5 |
| | BDI | 15.3 | 15 | 15.4 | 14.7 | 15.1 |
| Pollution index | PTI | 2.5 | 2.5 | 2.4 | 2.4 | 2.5 |
| | S | 2.3 | 2.3 | 2.4 | 2.4 | 2.4 |

composition of the stream, TDI, BDI, and chl-a values indicate that the stream is mesotrophic. So, diversity indices calculated using epilithic diatoms did not exactly represent the trophic status of the stream. Consequently, based on all these results, it can be said that Batlama stream does not yet reach the level of pollution and having mesotrophic characteristics as a trophic structure.

REFERENCES

- Pan Y, Stevenson RJ, Hill BH, Kaufmann PR, Herlihy AT. Spatial patterns and ecological determinants of benthic algal assemblages, stream slope, TP, TN and riparian canopy coverage. *J. of Phycology* 35 (1999) 460–468. DOI: 10.1046/j.1529-8817.1999.3530460.x
- Hill W, Dimick SM. Effects of riparian leaf dynamics on periphyton photosynthesis and light utilization efficiency. *Freshw. Biol.* 47 (2002) 1245–1256. DOI: 10.1046/j.1365-2427.2002.00837.x
- Lin HJ, Peng TR, Cheng IC, Chen LW, Kuo MH, Tzeng CS. A trophic model of the subtropical headwater stream habitat of the Formosan landlocked salmon, *Oncorhynchus formosanus*. *Aquatic Biol.* 17 (2012) 269–283. DOI: https://doi.org/10.3354/ab00481
- Quinn JM, Cooper AB, Stroud MJ, Burrell GP. Shade effects on stream periphyton and invertebrates: an experiment in streamside channels. *N. Z. J. of Mar. and Freshw.* 31 (1997) 665–683. DOI: 10.1080/00288330.1997.9516797
- Soininen J, Könönen K. Comparative study of monitoring South-Finnish rivers and streams using macroinvertebrate and benthic diatom community structure. *Aquatic Ecology* 38 (2004) 63–75. DOI:10.1023/B:AECO.0000021004.06965.bd
- Bowman MF, Chambers PA, Schindler DW. Epilithic algal abundance in relation to anthropogenic changes in phosphorus bioavailability and limitation in mountain rivers. *Canadian J. Fisheries Aquatic Sci.* 62(1) (2005) 174–184. DOI: 10.1139/f04-244
- Yıldız K. Investigations on algae communities of Meram stream, section 3–Algae living on sediment. *J. of Nat. Sci. A2*, 9(2) (1985) 428–434 (in Turkish).
- Kılınc S, Sivacı ER. A study on the past and present diatom flora of two alkaline lakes. *Turkish J. of Botany* 25 (2001) 373–378.
- Maraşlıoğlu F, Soylu EN, Gönüloğlu A. A study of the composition and seasonal variation of the epilithic diatoms of Lake Ladik (Samsun, Turkey). *International J. on Algae* 7(1) (2005) 58–70. DOI: 10.1615/InterJAlgae.v7.i1.40
- Maraşlıoğlu F, Soylu EN, Gönüloğlu A. Seasonal Variation and Occurrence of Epiphytic Diatom Assemblages on Mats of *Cladophora glomerata* in Lake Ladik, Samsun, Turkey. *Cryptogamie Algologie* 28(4) (2007) 373–384.
- Sivacı ER, Pabuçcu K. Epilithic and epilithic diatoms of the Balıklı Spain central Anatolia, Turkey. *J. of Freshw. Ecology* 22 (2007) 145–146. DOI: 10.1080/02705060.2007.9664154
- Soylu EN, Maraşlıoğlu F, Gönüloğlu A. Epilithic algae and seasonal variation of Gıcı Lake (Samsun–Bafra). *J. of Fisheries Sciences*. com 4(4) (2010) 437–445.
- Çiçek NL, Kalyoncu H., Akköz C., Ertan Ö. Seasonal Distribution and Species of Epilithic Algae in Darıören and Isparta Stream (Isparta–Turkey). *J. of Fisheries Sciences*. com 4(1) (2010) 78–90.
- Çolak-Sabancı F. Relationship of Epilithic Diatom Communities to Environmental Variables in Homa Lagoon (İzmir, Turkey). *Aquatic Biol.* 13 (2011) 233–241. DOI: 10.3354/ab00367
- Maraşlıoğlu F, Soylu EN, Gönüloğlu A. Seasonal Variations and Occurrence of Algal Turf Community in Lake Ladik (Turkey). *J. of Environmental Biol.* 34 (2013) 107–111.
- Ongun–Sevindik T, Küçük F. Benthic Diatoms as Indicators of Water Quality in the Acarlar Floodplain Forest (Northern Turkey). *Fresenius Environmental Bull.* 25(10) (2016) 4013–4025.
- Kalyoncu H, Lerzan–Çiçek N, Cengiz Akköz C, Yorulmaz B. Comparative performance of diatom indices in aquatic pollution assessment. *African J. of Agricultural Research* 4 (10) (2009) 1032–1040.
- Tokatlı C.. Use of Some Diatom Indices for Evaluating Water Quality: Sample of Gürleyik Stream (Eskişehir). *Dumlupınar University, J. of Nat. Sci.* 29 (2012) 21–28.
- Solak CN, Fehér G, Barlas M, Pabuçcu K. 2007. Use of epilithic diatoms to evaluate water quality of Akçay Stream (Büyük Menderes River) in Mugla/Turkey. *Large Rivers* 17 (3–4) (2007) 327–338. DOI: 10.1127/lr/17/2007/327
- Zelinka M, Marvan P. Zur Prazisierung derbiologischen Klassifikation des Reinheit fliessender Gewässer. *Arch. für Hydrobiologie* 57 (1961) 389–407.
- Round FE. An investigation of two benthic algal communities in Malharm Tarn, Yorkshire. *J. of Ecology* 41 (1953) 97–174. DOI: 10.2307/2257108
- Sladeckova A. Limnological investigation methods for the perifiton (Aufwuch) community. *Botanical Review* 28 (1962) 286–350.
- John DM, Whitton BA, Brook AJ. *The Freshwater Algal Flora of the British Isles: An identification guide to freshwater and terrestrial algae. The Natural History Museum and The British Phycological Society, Cambridge University Press,*

- Cambridge, 2003.
24. Krammer K, Lange-Bertalot H. Sübwasserflora von Mitteleuropa. Bacillariophyceae, 3. Teil. Centrales, Fragillariaceae, Eunoticeae. Gustav Fischer Verlag, Stuttgart, 1991a.
 25. Krammer K, Lange-Bertalot H. Sübwasserflora von Mitteleuropa. Bacillariophyceae, 4. Teil. Achnantheaceae, Kritische Ergänzungen zu Navicula (Lineolate) und Gomphonema Gesamtliterat. Gustav Fischer Verlag, Stuttgart, 1991b.
 26. Krammer K, Lange-Bertalot H. Sübwasserflora von Mitteleuropa. Bacillariophyceae, 1. Teil. Naviculaceae. Spectrum Acad. Verlag, Berlin, 1999a.
 27. Krammer K, Lange-Bertalot H. Sübwasserflora von Mitteleuropa. Bacillariophyceae, 2. Teil. Bacillariaceae, Epithemiaceae, Surirellaceae. Spectrum Academischer Verlag, Berlin, 1999b.
 28. Guiry MD, Guiry GM. AlgaeBase. World-wide electronic publication. Galway: National University of Ireland, 2016. Retrieved from <http://www.algaebase.org>.
 29. Brummitt RK, Powell CE. Authors of plant names. A list of authors of scientific names of plants, with recommended standart forms of their names, including abbreviations, Royal Botanic Gardens, Kew, 1992.
 30. Shannon CE, Weaver W. The Mathematical Theory of Communication. The University of Illinois Press, 1949.
 31. Kelly MG, Whitton BA. The trophic diatom index: A new index for monitoring eutrophication in rivers. J. of Appl. Phycology 7 (1995) 433-444. DOI: 10.1007/BF00003802
 32. Lenoir A, Coste M. Development of a practical diatom index of overall water quality applicable to the French National Water Board network, in: Whitton BA and Rott E (eds.) Use of Algae for Monitoring Rivers II. Institut für Botanik, Universität Innsbruck. pp. 29-43, 1996.
 33. Pantle R, Buck H. Die biologische Überwachung der Gewässer und die Darstellung der Ergebnisse (Biological monitoring of water bodies and the presentation of results). Gas -und Wasserfach 96 (1955) 604-618.
 34. Muscio C. The diatom pollution tolerance index: assigning tolerance values. Water Protection & Development Review Department. Environmental Resource Management, 2002.
 35. Strickland JDH, Parsons TR. A Pratical Handbook of Seawater Analysis. Fisheries Research Board of Canada, 1972.
 36. Sienkiewicz E. Limnological record inferred from diatoms in sediments of Lake Skaliska (north-eastern Poland). Acta Palaeobotanica 53(1) (2013) 99-104. DOI: 10.2478/acpa-2013-0007
 37. Gönüloğlu A, Öztürk M, Öztürk M. A Check-list of the Freshwater Algae of Turkey. Ondokuz Mayıs University, Faculty of Arts and Science, J. of Sci. 7(1) (1996) 8-46 (in Turkish).
 38. Gönüloğlu A. 2017. Turkishalgae Electronic Publication, Samsun, Turkey. <http://turkiyealgleri.omu.edu.tr>. (searched on 07 February 2017)
 39. Sivacı R, Dere Ş. Seasonal Changes of Epilithic (Aksaray-Ihlara) Diatom of Melendiz Stream and Effect on Total Organisms of Water Flow. Ekoloji 16(6) (2007) 29-36 (in Turkish).
 40. Yavuz O, Çetin K. Seasonal Variation and Algae on Pelagic Region of Cip Stream (Elazığ-Turkey). Firat University, J. of Sci. and Eng. 12(2) (2000) 25-39 (in Turkish).
 41. Wetzel RG. Limnology. Michigan State University. Academic Press, 2001.
 42. DSI. Water Quality and Dam Lakes which Provide Drinking Water of Ankara. DSI General Directory Research Reports, Ankara, 1998.
 43. Round FE. The Ecology of Algae. Cambridge University Press, Cambridge, 1981.
 44. Vinebrooke RD. Abiotic and Biotic Regulation of Periphyton in Recovering Acidified Lakes. J. of the N. Am. Benthological Soc. 15 (1996) 318-331. DOI: 10.2307/1467280
 45. Moses BS. The status of the fisheries downstream in river Niger the Kainji Dam, in: Proceedings of International Conference of Kainji Lake and River Basins Development in Africa. New Bussa, Nigeria, Kainji Lake Research Institute pp. 612, 1979.
 46. Köster D, Hübener T. Application of Diatom Indices in a planted ditch constructed for tertiary sewage treatment in Schwaan, Germany. Internationale Revue der Gesamten. Hidrobiologie 86 (2001) 241-252.